

## 6-6

## Reteaching

## Function Operations

When you combine functions using addition, subtraction, multiplication, or division, the domain of the resulting function has to include the domains of both of the original functions.

**Problem**

Let  $f(x) = x^2 - 4$  and  $g(x) = \sqrt{x}$ . What is the solution of each function operation? What is the domain of the result?

- $(f + g)(x) = f(x) + g(x) = (x^2 - 4) + (\sqrt{x}) = x^2 + \sqrt{x} - 4$
- $(f - g)(x) = f(x) - g(x) = (x^2 - 4) - (\sqrt{x}) = x^2 - \sqrt{x} - 4$
- $(g - f)(x) = g(x) - f(x) = (\sqrt{x}) - (x^2 - 4) = -x^2 + \sqrt{x} + 4$
- $(f \cdot g)(x) = f(x) \cdot g(x) = (x^2 - 4)(\sqrt{x}) = x^2\sqrt{x} - 4\sqrt{x}$

The domain of  $f$  is all real numbers. The domain of  $g$  is all  $x \geq 0$ . For parts a–d, there are no additional restrictions on the values for  $x$ , so the domain for each of these is  $x \geq 0$ .

$$e. \frac{f}{g}(x) = \frac{f(x)}{g(x)} = \frac{x^2 - 4}{\sqrt{x}} = \frac{(x^2 - 4)\sqrt{x}}{x}$$

As before, the domain is  $x \geq 0$ . But, because the denominator cannot be zero, eliminate any values of  $x$  for which  $g(x) = 0$ . The only value for which  $\sqrt{x} = 0$  is  $x = 0$ . Therefore, the domain of  $\frac{f}{g}$  is  $x > 0$ .

$$f. \frac{g}{f}(x) = \frac{g(x)}{f(x)} = \frac{\sqrt{x}}{x^2 - 4}$$

Similarly, begin with  $x \geq 0$  and eliminate any values of  $x$  that make the denominator  $f(x)$  zero:  $x^2 - 4 = 0$  when  $x = -2$  and  $x = 2$ . Therefore, the domain of  $\frac{g}{f}$  is  $x \geq 0$  combined with  $x \neq -2$  and  $x \neq 2$ . In other words, the domain is  $x \geq 0$  and  $x \neq 2$ , or all nonnegative numbers except 2.

**Exercises**

Let  $f(x) = 4x - 3$  and  $g(x) = x^2 + 2$ . Perform each function operation and then find the domain of the result.

1.  $(f + g)(x)$

2.  $(f - g)(x)$

3.  $(g - f)(x)$

4.  $(f \cdot g)(x)$

5.  $\frac{f}{g}(x)$

6.  $\frac{g}{f}(x)$

# 6-6


## Reteaching (continued)

### Function Operations

- One way to combine two functions is by forming a composite.
- A composite is written  $(g \circ f)$  or  $g(f(x))$ . The two different functions are  $g$  and  $f$ .
- Evaluate the inner function  $f(x)$  first.
- Use this value, the first output, as the input for the second function,  $g(x)$ .

#### Problem


What is the value of the expression  $g(f(2))$  given the inner function,  $f(x) = 3x - 5$  and the outer function,  $g(x) = x^2 + 2$ ?



#### Exercises

Evaluate the expression  $g(f(5))$  using the same functions for  $g$  and  $f$  as in the Example. Fill in blanks 7-14 on the chart.

Use one color highlighter to highlight the first input. Use a second color to highlight the first output and the second input. Use a third color to highlight the second output, which is the answer.



Given  $f(x) = x^2 + 4x$  and  $g(x) = 2x + 3$ , evaluate each expression.

15.  $f(g(2))$

16.  $g(f(2.5))$

17.  $g(f(-5))$

18.  $f(g(-5))$